arXiv:1610.07945v1 [math.NT] 25 Oct 2016

Real zeros of the Hurwitz zeta function

Toshiki Matsusaka

Abstract

It is well known that real zeros of the Riemann zeta function are negative even integers. As for the real zeros of the Hurwitz zeta function, T. Nakamura recently gave a non-vanishing condition in the intervals $(0,1)$ and $(-1,0)$. We generalize this result for all negative real numbers.

1 Introduction

The Hurwitz zeta function was first introduced by Hurwitz [2].

Definition 1.1. *Let $0 < a \leq 1$, $s \in \mathbb{C}$. Then the Hurwitz zeta function $\zeta(s,a)$ is defined by*

$$\zeta(s,a) := \sum_{n=0}^{\infty} (n+a)^{-s}, \quad s := \sigma + it, \quad \sigma > 1, \quad t \in \mathbb{R}.$$

This series converges absolutely in the half-plane $\sigma > 1$ and uniformly in each compact subset of this half-plane. Moreover $\zeta(s,a)$ is analytically continued to the whole complex plane except for a simple pole at $s = 1$. We easily see that the Hurwitz zeta function has no real zeros in this half-plane $\sigma > 1$. For the special case of $a = 1$, the Riemann zeta function $\zeta(s) = \zeta(s,1)$ has real zeros at the negative even integers called the trivial zeros. In general, Nakamura [4, 5] investigated real zeros of $\zeta(s,a)$ in the intervals $(0,1)$ and $(-1,0)$, and obtained the following non-vanishing conditions.

Theorem 1.2. *([4, 5]) Let $b_2^{\pm} := (3 \pm \sqrt{3})/6$. Then*

(1) *$\zeta(\sigma,a)$ does not vanish when $\sigma \in (0,1)$ if and only if $1/2 \leq a \leq 1$,*

(2) *$\zeta(\sigma,a)$ does not vanish when $\sigma \in (-1,0)$ if and only if $b_2^- \leq a \leq 1/2$ or $b_2^+ \leq a \leq 1$.*

Note that $b_2^{\pm}$ are the roots of the second Bernoulli polynomial $B_2(x) := x^2 - x + 1/6$.

In this article, we generalize this result to general intervals in the negative real numbers.

Theorem 1.3. *Let $N \geq 0$ be an integer. Then $\zeta(\sigma,a)$ does not vanish when $\sigma \in (-N-1,-N)$ if and only if*

$$\begin{cases} b_{N+2}^- \leq a \leq 1/2 \text{ or } b_{N+2}^+ \leq a \leq 1 & \text{if } N \text{ is even,} \\ 0 < a \leq b_{N+1}^- \text{ or } 1/2 \leq a \leq b_{N+1}^+ & \text{if } N \text{ is odd,} \end{cases}$$

where $b_n^{\pm}$ are the two roots of the nth Bernoulli polynomial $B_n(x)$ in the interval $(0,1)$ (defined in Section 2).

Including the case of $N = -1$, we can also state Theorem 1.3 in the following form.

Theorem 1.4. *Let $N \geq -1$ be an integer. Then $\zeta(\sigma, a)$ does not vanish when $\sigma \in (-N-1, -N)$ if and only if $B_{N+1}(a)B_{N+2}(a) \geq 0$.*

We can easily see that Theorem 1.2 is a special case of Theorem 1.4. In Section 2, we show some properties of the Bernoulli polynomials and the Hurwitz zeta function. In Section 3, we prove Theorem 1.4 for the cases of $1 \leq N \leq 3$ and $N \geq 4$, separately. Note that the known results on zeros of the Hurwitz zeta function are reviewed in Nakamura [5, Section 1.2].

2 Preliminaries

In this section, we recall some properties of the Bernoulli polynomials and the Hurwitz zeta function. First, we define the Bernoulli numbers and the Bernoulli polynomials.

Definition 2.1. (*see* [*1*]) *The Bernoulli numbers B_n and the Bernoulli polynomials $B_n(x)$ are defined by means of generating functions,*

$$\begin{aligned} \sum_{n=0}^{\infty} B_n \frac{t^n}{n!} &= \frac{te^t}{e^t - 1}, \\ \sum_{n=0}^{\infty} B_n(x) \frac{t^n}{n!} &= \frac{te^{xt}}{e^t - 1}. \end{aligned}$$

We can easily see that $B_n(1) = B_n$ holds.

It is known that the nth Bernoulli polynomial $B_n(x)$ has only one root in each interval $[0, 1/2)$ and $[1/2, 1)$ for $n \geq 2$. (see [3, Section 1]). Then, we can define b_n^- and b_n^+ as the roots of $B_n(x)$ in the intervals $[0, 1/2)$ and $[1/2, 1)$, respectively. For further properties of the Bernoulli polynomials, see [1, 3]. Next, we show an integral representation of the Hurwitz zeta function.

Proposition 2.2. *Let $N \geq -1$ be an integer. For $-N - 1 < \mathrm{Re}(s) < -N$, it holds that*

$$\Gamma(s)\zeta(s, a) = \int_0^{\infty} \left(\frac{e^{(1-a)x}}{e^x - 1} - \sum_{n=0}^{N+1} \frac{B_n(1-a)}{n!} x^{n-1} \right) x^{s-1} dx.$$

Proof. It is known that the Hurwitz zeta function has the integral representation

$$\zeta(s, a) = \frac{1}{\Gamma(s)} \int_0^{\infty} \frac{e^{(1-a)x}}{e^x - 1} x^{s-1} dx, \quad \mathrm{Re}(s) > 1,$$

(see [1]). Note that the integrand function can be expanded as the power series

$$\frac{e^{(1-a)x}}{e^x - 1} = \sum_{n=0}^{\infty} \frac{B_n(1-a)}{n!} x^{n-1}$$

for $|x| < 2\pi$.

First, we meromorphically continue the function $\Gamma(s)\zeta(s,a)$ to the half-plane $\mathrm{Re}(s) > -N-1$ by using this integral representation. We divide this integral into three parts,

$$\begin{aligned}
\Gamma(s)\zeta(s,a) &= P(s) + Q_N(s) + R_N(s), \\
P(s) &= \int_1^\infty \frac{e^{(1-a)x}}{e^x - 1} x^{s-1} dx, \\
Q_N(s) &= \int_0^1 \Big(\sum_{n=0}^{N+1} \frac{B_n(1-a)}{n!} x^{n-1} \Big) x^{s-1} dx, \\
R_N(s) &= \int_0^1 \Big(\frac{e^{(1-a)x}}{e^x - 1} - \sum_{n=0}^{N+1} \frac{B_n(1-a)}{n!} x^{n-1} \Big) x^{s-1} dx.
\end{aligned}$$

Then $P(s)$ is holomorphic in the whole complex plane, $Q_N(s)$ is a rational function, and $R_N(s)$ is holomorphic in the half-plane $\mathrm{Re}(s) > -N-1$, that is, $\Gamma(s)\zeta(s,a)$ is meromorphically continued to the half-plane $\mathrm{Re}(s) > -N-1$.

Next we restrict the domain to the strip $-N-1 < \mathrm{Re}(s) < -N$. Then we can write

$$\begin{aligned}
Q_N(s) &= \sum_{n=0}^{N+1} \frac{B_n(1-a)}{n!} \frac{1}{n+s-1} \\
&= -\sum_{n=0}^{N+1} \frac{B_n(1-a)}{n!} \int_1^\infty x^{n+s-2} dx \\
&= -\int_1^\infty \sum_{n=0}^{N+1} \frac{B_n(1-a)}{n!} x^{n+s-2} dx,
\end{aligned}$$

the last integral being absolutely convergent. We therefore obtain

$$P(s) + Q_N(s) + R_N(s) = \int_0^\infty \Big(\frac{e^{(1-a)x}}{e^x - 1} - \sum_{n=0}^{N+1} \frac{B_n(1-a)}{n!} x^{n-1} \Big) x^{s-1} dx$$

for $-N-1 < \mathrm{Re}(s) < -N$. □

Finally, we remark the classical relation between the Bernoulli polynomial and the Hurwitz zeta function.

Lemma 2.3. (*See [1]*) *For each integer $n > 0$, it holds that*

$$\zeta(1-n, a) = -\frac{B_n(a)}{n}.$$

3 Proof of Theorem 1.4

We give a proof for the cases of $1 \le N \le 3$ and $N \ge 4$.

(1) $1 \le N \le 3$ case.

For each integer $N \geq -1$, We put

$$G_N(a,x) := \frac{e^{(1-a)x}}{e^x - 1} - \sum_{n=0}^{N+1} \frac{B_n(1-a)}{n!} x^{n-1}.$$

Then we have $\Gamma(s)\zeta(s,a) = \int_0^\infty G_N(a,x)x^{s-1}dx$ for $-N-1 < \mathrm{Re}(s) < -N$ by Proposition 2.2. Moreover we put

$$g_N(a,x) := x(e^x-1)G_N(a,x) = xe^{(1-a)x} - \sum_{n=0}^{N+1} \frac{B_n(1-a)}{n!} x^n(e^x-1),$$

and differentiate it $N+2$ times. Then we get $g_N^{(k)}(a,0) = 0$ for $k = 0, 1, 2, \ldots, N+2$. By direct calculation, we have

$$\begin{aligned} g_N^{(N+2)}(a,x) &= (N+2)(1-a)^{N+1}e^{(1-a)x} + (1-a)^{N+2}xe^{(1-a)x} \\ &\quad - \sum_{n=0}^{N+1} \frac{B_n(1-a)}{n!} \sum_{k=0}^{n} \binom{N+2}{k}\binom{n}{k} k! x^{n-k} e^x. \end{aligned}$$

Putting $m = n-k$, we can get the form

$$\begin{aligned} e^{(a-1)x} g_N^{(N+2)}(a,x) &= (N+2)(1-a)^{N+1} + (1-a)^{N+2}x \\ &\quad - \sum_{m=0}^{N+1} \left(\sum_{k=0}^{N+1-m} \binom{N+2}{k} B_{m+k}(1-a) \right) \frac{x^m}{m!} \cdot e^{ax}. \end{aligned}$$

(1-i) For the case of $N = 1$, we have

$$\begin{aligned} & e^{(a-1)x} g_1^{(3)}(a,x) \\ &= 3(1-a)^2 + (1-a)^3 x - \left(3(1-a)^2 + (1-3a)(1-a)x + \frac{1}{2}B_2(1-a)x^2 \right) \cdot e^{ax} \\ &= -2a(1-a)(1-2a)x \\ &\qquad + \sum_{n=2}^{\infty} a^{n-2} \left(-3(1-a)^2a^2 - n(1-3a)(1-a)a - \frac{n(n-1)}{2}B_2(1-a) \right) \frac{x^n}{n!}. \end{aligned}$$

If $0 < a \leq b_2^-$, we have $-2a(1-a)(1-2a) < 0$. For $n \geq 2$, we have

$$\begin{aligned} -3(1-a)^2a^2 &< 0, \\ -n(1-3a)(1-a)a &< 0, \\ -\frac{n(n-1)}{2}B_2(1-a) &\leq 0, \end{aligned}$$

that is, $e^{(a-1)x}g_1^{(3)}(a,x) < 0$ for all $x > 0$. By $g_1(a,0) = g_1^{(1)}(a,0) = g_1^{(2)}(a,0) = g_1^{(3)}(a,0) = 0$, we obtain $g_1(a,x) < 0$, $G_1(a,x) < 0$, and thus $\zeta(\sigma,a) < 0$ for all $\sigma \in (-2,-1)$.

If $1/2 \leq a \leq b_2^+$, we have $-2a(1-a)(1-2a) \geq 0$. For $n \geq 2$, we have

$$\begin{aligned} -n(1-3a)(1-a)a &> 0, \\ -\frac{n(n-1)}{2}B_2(1-a) &\geq 0. \end{aligned}$$

For the remaining part, it holds that

$$\begin{aligned} -3(1-a)^2a^2 - n(1-3a)(1-a)a &= a(1-a)\{-3(1-a)a - n(1-3a)\} \\ &\geq a(1-a)\{-3(1-a)a - 2(1-3a)\} \\ &= -a(1-a)(2-3a-3a^2) > 0. \end{aligned}$$

Thus we obtain $e^{(a-1)x}g_1^{(3)}(a,x) > 0$, that is, $\zeta(\sigma,a) > 0$ for all $\sigma \in (-2,-1)$ in the same way.

Finally if $b_2^- < a < 1/2$ or $b_2^+ < a \leq 1$, then it holds that $B_2(a)B_3(a) < 0$. By Lemma 2.3, we see that $\zeta(-1,a) = -B_2(a)/2$ and $\zeta(-2,a) = -B_3(a)/3$. By assumption, we have $\zeta(-1,a)\zeta(-2,a) < 0$, that is, $\zeta(\sigma,a)$ has zeros in $(-2,-1)$. This concludes the proof for the case of N =1.

(1-ii) For the case of $N = 2$, we have

$$\begin{aligned} &e^{(a-1)x}g_2^{(4)}(a,x) \\ &\quad = -\frac{1}{6}(1-30a^2+60a^3-30a^4)x - \frac{1}{12}(1-4a-30a^2+60a^3-24a^5)x^2 \\ &\qquad + \sum_{n=3}^{\infty} a^{n-3}\left(-4(1-a)^3a^3 - \frac{n}{6}(7-6a)(1-6a+6a^2)a^2\right. \\ &\qquad \left. -\frac{n(n-1)}{12}(1-18a+42a^2-24a^3)a - \frac{n(n-1)(n-2)}{6}B_3(1-a)\right)\frac{x^n}{n!}. \end{aligned}$$

If $b_4^- \leq a \leq 1/2$, we have

$$\begin{aligned} -\frac{1}{6}(1-30a^2+60a^3-30a^4) &\geq 0, \\ -\frac{1}{12}(1-4a-30a^2+60a^3-24a^5) &> 0. \end{aligned}$$

For $n \geq 3$, we have

$$\begin{aligned} -\frac{n}{6}(7-6a)(1-6a+6a^2)a^2 &> 0, \\ -\frac{n(n-1)}{12}(1-18a+42a^2-24a^3)a &> 0, \\ -\frac{n(n-1)(n-2)}{6}B_3(1-a) &\geq 0. \end{aligned}$$

For the remaining part, it holds that

$$\begin{aligned} &-4(1-a)^3a^3 - \frac{n(n-1)}{12}(1-18a+42a^2-24a^3)a \\ &\geq -4(1-a)^3a^3 - \frac{3(3-1)}{12}(1-18a+42a^2-24a^3)a \\ &= -\frac{1}{2}(1-18a+50a^2-48a^3+24a^4-8a^5)a > 0. \end{aligned}$$

Thus we obtain $e^{(a-1)x}g_2^{(4)}(a,x) > 0$ for all $x > 0$, that is, $\zeta(\sigma,a) < 0$ for all $\sigma \in (-3,-2)$.

If $b_4^+ \leq a \leq 1$, we have

$$\begin{aligned} -\frac{1}{6}(1-30a^2+60a^3-30a^4) &\leq 0, \\ -\frac{1}{12}(1-4a-30a^2+60a^3-24a^5) &< 0. \end{aligned}$$

For $n \geq 3$, we have

$$\begin{aligned} -4(1-a)^3a^3 &\leq 0, \\ -\frac{n(n-1)}{12}(1-18a+42a^2-24a^3)a &< 0, \\ -\frac{n(n-1)(n-2)}{6}B_3(1-a) &\leq 0. \end{aligned}$$

For the remaining part, it holds that

$$\begin{aligned} &-\frac{n}{6}(7-6a)(1-6a+6a^2)a^2 - \frac{n(n-1)}{12}(1-18a+42a^2-24a^3)a \\ &\leq n\left\{-\frac{1}{6}(7-6a)(1-6a+6a^2)a^2 - \frac{(3-1)}{12}(1-18a+42a^2-24a^3)a\right\} \\ &= -\frac{n}{6}(1-11a-6a^2+54a^3-36a^4)a < 0. \end{aligned}$$

Thus we obtain $e^{(a-1)x}g_2^{(4)}(a,x) < 0$, that is, $\zeta(\sigma,a) > 0$ for all $\sigma \in (-3,-2)$.

Finally if $0 < a < b_4^-$ or $1/2 < a < b_4^+$, then $\zeta(\sigma,a)$ has zeros in $(-3,-2)$ in the same way as (1-i). This concludes the proof for the case of N =2.

(1-iii) For the case of $N = 3$, we have

$$\begin{aligned} &e^{(a-1)x}g_3^{(5)}(a,x) \\ &= a(1-a)(1-2a)(1+3a-3a^2)x + \frac{1}{12}(1+9a-9a^2-90a^3+120a^4-30a^6)x^2 \\ &\quad + \frac{1}{36}(1+6a+6a^2-99a^3+180a^5-60a^6-30a^7)x^3 \\ &\quad + \sum_{n=4}^{\infty} a^{n-4}\Bigg(-5(1-a)^4a^4 - n(1-a)(1-10a+20a^2-10a^3)a^3 \\ &\quad + \frac{n(n-1)}{12}(1+21a-111a^2+150a^3-60a^4)a^2 \\ &\quad + \frac{n(n-1)(n-2)}{36}(1+3a-39a^2+66a^3-30a^4)a - \frac{n(n-1)(n-2)(n-3)}{24}B_4(1-a)\Bigg)\frac{x^n}{n!}. \end{aligned}$$

If $0 < a \leq b_4^-$, we have

$$\begin{aligned} a(1-a)(1-2a)(1+3a-3a^2) &> 0, \\ \frac{1}{12}(1+9a-9a^2-90a^3+120a^4-30a^6) &> 0, \\ \frac{1}{36}(1+6a+6a^2-99a^3+180a^5-60a^6-30a^7) &> 0. \end{aligned}$$

For $n \geq 4$, we have

$$\begin{aligned} \frac{n(n-1)}{12}(1+21a-111a^2+150a^3-60a^4)a^2 &> 0, \\ \frac{n(n-1)(n-2)}{36}(1+3a-39a^2+66a^3-30a^4)a &> 0, \\ -\frac{n(n-1)(n-2)(n-3)}{24}B_4(1-a) &\geq 0. \end{aligned}$$

For the remaining parts, we have

$$\begin{aligned}
&-5(1-a)^4a^4+\frac{n(n-1)}{12}(1+21a-111a^2+150a^3-60a^4)a^2\\
&\geq -5(1-a)^4a^4+\frac{4(4-1)}{12}(1+21a-111a^2+150a^3-60a^4)a^2\\
&=(1+21a-116a^2+170a^3-90a^4+20a^5-5a^6)a^2>0,
\end{aligned}$$

and

$$\begin{aligned}
&-n(1-a)(1-10a+20a^2-10a^3)a^3+\frac{n(n-1)(n-2)}{36}(1+3a-39a^2+66a^3-30a^4)a\\
&\geq n\left\{-(1-a)(1-10a+20a^2-10a^3)a^3+\frac{(4-1)(4-2)}{36}(1+3a-39a^2+66a^3-30a^4)a\right\}\\
&=\frac{n}{6}(1+3a-45a^2+132a^3-210a^4+180a^5-60a^6)a>0.
\end{aligned}$$

Thus we obtain $e^{(a-1)x}g_3^{(5)}(a,x)>0$ for all $x>0$, that is, $\zeta(\sigma,a)>0$ for all $\sigma\in(-4,-3)$.

If $1/2\leq a\leq b_4^+$, we have

$$\begin{aligned}
a(1-a)(1-2a)(1+3a-3a^2) &\leq 0,\\
\frac{1}{12}(1+9a-9a^2-90a^3+120a^4-30a^6) &< 0,\\
\frac{1}{36}(1+6a+6a^2-99a^3+180a^5-60a^6-30a^7) &< 0.
\end{aligned}$$

For $n\geq 4$, we have

$$\begin{aligned}
-5(1-a)^4a^4 &< 0,\\
\frac{n(n-1)}{12}(1+21a-111a^2+150a^3-60a^4)a^2 &< 0,\\
\frac{n(n-1)(n-2)}{36}(1+3a-39a^2+66a^3-30a^4)a &< 0,\\
-\frac{n(n-1)(n-2)(n-3)}{24}B_4(1-a) &\leq 0.
\end{aligned}$$

For the remaining part, it holds that

$$\begin{aligned}
&-n(1-a)(1-10a+20a^2-10a^3)a^3+\frac{n(n-1)}{12}(1+21a-111a^2+150a^3-60a^4)a^2\\
&\leq n\left\{-(1-a)(1-10a+20a^2-10a^3)a^3+\frac{(4-1)}{12}(1+21a-111a^2+150a^3-60a^4)a^2\right\}\\
&=\frac{n}{4}(1+17a-67a^2+30a^3+60a^4-40a^5)a^2<0.
\end{aligned}$$

Thus we obtain $e^{(a-1)x}g_3^{(5)}(a,x)<0$, that is, $\zeta(\sigma,a)<0$ for all $\sigma\in(-4,-3)$.

Finally if $b_4^-<a<1/2$ or $b_4^+<a\leq 1$, then $\zeta(\sigma,a)$ has zeros in $(-4,-3)$. This concludes the proof for the case of N =3.

(2) $N\geq 4$ case.

We give a proof of the remaining cases of $N\geq 4$ by using Spira's theorem.

Theorem 3.1. ([6]) *If $\sigma \leq -(4a+1+2[1-2a])$ and $|t| \leq 1$, then $\zeta(s,a) \neq 0$ except for zeros on the negative real axis, one in each interval $(-2n-4a-1, -2n-4a+1)$ for $n \geq 1-2a$.*

Corollary 3.2. *Let $M \geq 2$ be an integer. The Hurwitz zeta function $\zeta(\sigma, a)$ has only one zero in each interval $[-2M-2, -2M)$.*

Proof. By definition, we have $\zeta(s,1/2) = (2^s-1)\zeta(s)$ and $\zeta(s,1) = \zeta(s)$. Thus, for $a = 1/2$ and 1, the point $s = -2M-2$ is the only one zero in each $[-2M-2, -2M)$. In other cases, the Hurwitz zeta function has an odd number of real zeros counting multiplicity in each $[-2M-2, -2M)$ by $B_{2M+1}(a)B_{2M+3}(a) < 0$ and Lemma 2.3. However, by Theorem 3.1, the Hurwitz zeta function does not have more than two real zeros in each interval $[2M-2, -2M)$ whose length is 2. □

Theorem 1.4 for the cases of $N \geq 4$ follows readily from this corollary and Lemma 2.3. Combining (1-i, ii, iii) and Theorem 1.2, we therefore obtain Theorem 1.4 for the all cases of $N \geq -1$.

Acknowledgement. *The author thanks Professor Masanobu Kaneko and Professor Takashi Nakamura for their helpful comments on an earlier version of this paper.*

References

[1] T. Arakawa, T. Ibukiyama, M. Kaneko, Bernoulli Numbers and Zeta Functions, Springer Monographs in Mathematics, Springer, Tokyo, 2014, with an appendix by Don Zagier.

[2] A. Hurwitz, Einige Eigenschaften der Dirichletschen Functionen $F(s) = \sum(\frac{D}{n}) \cdot \frac{1}{n^s}$ die bei der Bestimmung der Klassenzahlen binärer quadratischer Formen auftreten, Z. Math. Phys., 27 (1882) 86–101.

[3] D. J. Leeming, The real zeros of the Bernoulli polynomials, Journal of Approximation theory, 58 (1989) 124–150.

[4] T. Nakamura, Real zeros of Hurwitz-Lerch zeta and Hurwitz-Lerch type of Euler-Zagier double zeta functions, Math. Proc. Cambridge Philos. Soc., 160 (1) (2016) 39–50.

[5] T. Nakamura, Real zeros of Hurwitz-Lerch zeta functions in the interval $(-1, 0)$, J. Math. Anal. Appl., 438 (1) (2016) 42–52.

[6] R. Spira, Zeros of Hurwitz zeta functions, Mathematics of computation, 30 (136) (1976) 863–866.

T. Matsusaka: Graduate School of Mathematics, Kyushu University, Motooka 744, Nishi-ku Fukuoka 819-0395, Japan
e-mail: toshikimatsusaka@gmail.com